


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/jasi

Original Article

Pre and postganglionic innervation of rat adrenal gland by fluorescent tract tracer – Fast blue[☆]

Santosh K. Sangari^{a,*}, Kamlesh Khatri^b

^a Program in Anatomy and Body Visualization, Department of Cell and Developmental Biology, Weill Cornell Medical College, 1300 York Avenue, NY 10065, USA

^b Department of Anatomy, School of Medical Sciences and Research, Sharda University, Greater Noida, Uttar Pradesh, India

ARTICLE INFO

Article history:

Received 25 March 2014

Accepted 16 April 2014

Available online 10 May 2014

Keywords:

Albino rat

Retrograde labeling

Sympathetic nerve supply

Suprarenal gland

ABSTRACT

Introduction: To search for pre and postganglionic neurons innervating the adrenal gland by injecting retrograde tract tracer fast blue in the adrenal medulla.

Methods: The motor innervation of rat adrenal gland was studied by a fluorescent tract tracer fast blue. 5 μ l of 2% aqueous suspension of fast blue was injected into left adrenal gland. After a survival period of 4–5 days, spinal cord, sympathetic ganglia, suprarenal ganglion, coeliac ganglion and left adrenal gland were dissected out and 15 μ m thick plastic sections (JB4 Polysciences) were examined under a fluorescent microscope.

Results: Retrogradely labeled preganglionic neurons were observed in the ipsilateral intermediolateral column of spinal cord from T3 to L2 spinal segments with maximum concentration of labeled neurons from T6 to T11. The labeled neurons were multipolar, spherical or fusiform in shape with transverse diameter 10–20 μ m and vertical diameter varying from 12 to 30 μ m. Postganglionic labeled neurons were also observed in the left suprarenal ganglion and left sympathetic ganglia (T5–L2) with maximum concentration from T6 to L1. Labeled neurons varied from 12 to 30 μ m in diameter and were randomly distributed throughout the ganglion.

Discussion: The preganglionic neurons from T3 to L2 spinal segments and postganglionic nerve fibers from ipsilateral sympathetic ganglia (T5–L2) and suprarenal ganglion supplying the adrenal gland might be responsible for the hormone release by regulating blood flow and also by directly innervating the parenchymal cells.

Copyright © 2014, Anatomical Society of India. Published by Reed Elsevier India Pvt. Ltd. All rights reserved.

[☆] Institute where work was conducted: Department of Anatomy, University College of Medical Sciences, Delhi 110095, India.

* Corresponding author. Tel.: +1 7325938021.

E-mail address: sks2005@med.cornell.edu (S.K. Sangari).

<http://dx.doi.org/10.1016/j.jasi.2014.04.008>

0003-2778/Copyright © 2014, Anatomical Society of India. Published by Reed Elsevier India Pvt. Ltd. All rights reserved.

1. Introduction

It is generally accepted that chromaffin cells of adrenal medulla are supplied by preganglionic cholinergic fibers, the cell bodies of which are located in the intermediolateral column of the spinal cord. Noxious chemical stimulation of thoracic and lumbar interspinous tissue in anesthetized rats is associated with large increase in adrenal sympathetic efferent nerve activity and catecholamine secretion.¹

The review of literature shows lot of variation regarding the extent of intermediolateral column supplying the adrenal gland.^{2–4}

The presence of enkephalin, substance P and vasoactive intestinal polypeptide like immunoreactivity has been seen in the nerve terminals of human adrenal medulla, which suggests that innervation to adrenal medulla may not be purely cholinergic.^{5–7} Acetylcholinesterase positive nerve plexus have also been demonstrated in the human adrenal cortex.^{8,9}

Adrenergic innervation has been demonstrated in cat adrenal medulla,¹⁰ whereas the postganglionic nerve endings have also been seen in the adrenal gland of dogfish and rat.^{11,12}

Therefore the present work was planned to search for postganglionic neurons innervating the adrenal gland and to confirm the extent of intermediolateral column of spinal cord supplying it by injecting retrograde tract tracer fast blue in the adrenal medulla.

2. Materials and methods

The study was conducted on 25 adult albino rats of either sex weighing between 200 and 250 g. The animals were kept under standard laboratory conditions with food and water ad libitum. Under ether anesthesia, laparotomy was performed and left adrenal gland was exposed. 5 μ l of 2% aqueous solution of fast blue was slowly injected over a period of 15 min. The needle of the microsyringe was kept in the gland for 5 min and then slowly withdrawn after the completion of injection to allow the tracer to pass from the syringe into the gland. The intestine was replaced in position and the anterior abdominal wall was sutured. After a survival period of 4–5 days, the rat was perfused with 10% formaldehyde. Anterior abdominal wall was opened. Left adrenal gland, celiac ganglion, suprarenal ganglion and C8 to L4 sympathetic ganglia of both sides were dissected out. Laminectomy was performed to expose the spinal cord. The spinal cord was taken out from C7 to L4 spinal segments.

The tissues were immersed in 10% sucrose cacodylate buffer for 24 h and embedded in plastic resin (JB4 kit polysciences) without prior dehydration. The 15 μ m thick serial sections of the spinal cord (longitudinal and transverse), sympathetic ganglia, suprarenal ganglion and celiac ganglion were examined under a fluorescent microscope using an excitation filter of 365 nm.

3. Results

Screening of sections of the left adrenal gland showed a needle track with maximum concentration of fluorescent dye in

the medulla of the adrenal gland. In three animals, the leakage was seen in the connective tissue outside the adrenal gland and were discarded from the study.

Retrogradely labeled neurons were seen in the intermediolateral column of the spinal cord from third thoracic to second lumbar spinal segments. Labeling was exclusively ipsilateral to the site of injection and no labeled neurons were seen on the opposite side.

Maximum concentration of labeled neurons was observed from T6 to T11 spinal segments. The labeled neurons were multipolar and showed bright blue fluorescence in the cytoplasm with a negative nuclear shadow. Fig. 1 shows transverse section of the spinal cord (T9 segment) with a labeled neuron in the left intermediolateral column of spinal cord. The labeled neurons were spherical or fusiform in shape with transverse diameter varying from 10 μ m to 20 μ m.

Longitudinal section of the spinal cord showed vertical disposition of the labeled neurons with bright blue fluorescence in the cytoplasm and a negative nuclear shadow (Fig. 2). The neurons vary from 20 μ m to 30 μ m in length.

Retrogradely labeled neurons were also observed in the suprarenal ganglion and from fifth thoracic to second lumbar sympathetic ganglia with maximum concentration from T6 to T11. Labeling was ipsilateral to the site of injection. Labeled neurons were 12 μ m–30 μ m in diameter and were arranged singly or in groups of 2–3 neurons throughout the ganglion without any specific localization (Fig. 3). No labeling was seen in the celiac ganglion.

4. Discussion

In the present study, fluorescent tract tracer fast blue was picked up by the nerve terminals in the adrenal medulla and the adjoining part of the adrenal cortex and labeled the neurons retrogradely in the spinal cord, sympathetic ganglia and suprarenal ganglion. These observations clearly demonstrate that the adrenal gland is supplied by both pre and post-ganglionic sympathetic nerve fibers.

It has been observed that the unilateral destruction of the adrenal medulla or the nerve terminals supplying the adrenal medulla leads to disappearance of preganglionic neurons of the intermediolateral column of T7 to T10 spinal




Fig. 1 – Transverse section of the spinal cord (T9) showing a labeled neuron in the left intermediolateral column of spinal cord. $\times 300$.



Fig. 2 – Longitudinal section of the spinal cord (T9–T10) showing vertically disposed labeled neurons. $\times 300$.

segments.^{13,14} But it is remarkable that in the present study, retrogradely labeled preganglionic neurons were observed in the vast extent of the intermediolateral column of the spinal cord from T3 to L2 spinal segments. The maximum concentration of the labeled neurons was observed from T6 to T11 spinal segments, which corresponds with the spinal segments contributing to splanchnic nerves. Our findings agree well with the reports available that the adrenal gland of rat, guinea pig and marmoset is supplied by preganglionic sympathetic fibers from intermediolateral column of T3 to L2 spinal segments.^{11,15} Whereas other studies observed that the intermediolateral column of T5–T13,¹⁶ C8 to T13,¹⁷ T2–T13² and T1 to L1⁴ spinal segments innervated rat adrenal gland by horseradish peroxidase and fluoro gold.

The extent of preganglionic neurons in the intermediolateral column of spinal cord supplying the adrenal gland in the present study also corresponds with the labeling of the adrenal sensory neurons in the dorsal root ganglia.^{18–20}

Labeling in the spinal cord was ipsilateral to the site of injection, which suggests that the innervation to the adrenal gland is exclusively ipsilateral with no crossing over of the nerve fibers to the opposite side. It also excludes the possibility of generalized vascular spread of the dye from the




Fig. 3 – Transverse section of left sympathetic ganglion (T10). Arrow points to the labeled sympathetic neuron $\times 300$.

injection site, as otherwise it would have produced bilateral labeling. An injection of Rhodamine Dextran Lysine into adrenal medulla has also shown ipsilateral labeling of spinal cord from T3 to T12 with distinct labeling from T7 to T10 spinal segments.²¹

The present retrograde tract tracing technique shows that in addition to sympathetic preganglionic neurons in the intermediolateral column of the spinal cord, postganglionic neurons in the sympathetic ganglia and suprarenal ganglion also innervate the adrenal gland.

The random distribution of labeled neurons in the suprarenal and sympathetic ganglia probably suggests that there is no viscerotopic representation of the adrenal gland in suprarenal and sympathetic ganglia. Similar observations were also made in the dorsal root ganglia for the organization of visceral sensory neurons of cat²² and for innervation of rat stomach.²³

No labeling was seen in celiac ganglion in the present study. This probably gives conclusive evidence that in rat, postganglionic neurons of celiac ganglion do not project to the adrenal gland.¹¹

Labeling was observed from T5 to L2 ipsilateral sympathetic ganglia with maximum concentration of labeled neurons from T6 to T11. These observations show that the postganglionic innervation from sympathetic ganglia closely corresponds to the preganglionic sympathetic outflow to the adrenal gland.

Electrophysiological evoked sympathetic responses have demonstrated the presence of both pre and postganglionic sympathetic fibers in greater splanchnic nerve²⁴ and in the adrenal cortex and medulla.²⁵ The presence of adrenergic fibers have been localized in the adrenal medulla of cat, distinctly and significantly associated with norepinephrine containing cell areas.¹⁰

The presence of VIP immunoreactive fibers in the adrenal cortex and medulla suggest that the innervation of adrenal cortex is involved in the adrenocortical rhythm of hormone release and stimulation of these nerves could concomitantly release hormones and increase the blood flow to the adrenal medulla.²⁶

Conflicts of interest

All authors have none to declare.

REFERENCES

1. Bolton P, Budgell B, Kimpton A. Influence of innocuous cervical vertebral movement on the efferent innervation of the adrenal gland in the rat. *Aut Neurosci : Basic Clin.* 2006;124(1):103–111.
2. Haase P, Contestabile A, Flumerfelt BA. Preganglionic innervation of the adrenal gland of the rat using horseradish peroxidase. *Exp Neurol.* 1982;78:217–221.
3. Holets V, Elde R. The differential distribution and relationship of serotonergic and peptidergic fibres to sympathoadrenal neurons in the intermediolateral cell column of the rat: a combined retrograde axonal transport and

- immunofluorescence study. *Neuroscience*. 1982;7(5):1155–1174.
4. Schramm LP, Adair JR, Stribling JM, Gray LP. Preganglionic innervation of the adrenal gland of the rat: a study using horseradish peroxidase. *Comp Neurol*. 1975;49:540–553.
 5. Berghorn KA, Li C, Nathanielsz PW, McDonald TJ. VIP innervation: sharp contrast in fetal sheep and baboon adrenal glands suggest differences in developmental regulation. *Brain Res*. 2000;877(2):271–280.
 6. Linnoila RJ, Diagustine RP, Hervonen A, Miller RJ. Distribution of (Met 5) - and (Leu 5) - enkephalin vasoactive intestinal polypeptide and substance P-like immunoreactivity in human adrenal glands. *Neuroscience*. 1980;5:2247–2259.
 7. Schultzberg M, Lundberg JM, Hökfelt T, et al. Enkephalin – like immunoreactivity in gland cells and nerve terminals of the adrenal medulla. *Neuroscience*. 1978;3:1169–1186.
 8. Janossy A, Orso E, Szalay KS, et al. Cholinergic regulation of the rat adrenal zona glomerulosa. *J Endocrinol*. 1998;157(2):305–315.
 9. Charlton BG, Nkomazana OF, McGadey J, Neal DE. A preliminary study of acetylcholinesterase- positive innervation in the human adrenal cortex. *J Anat*. 1991;176:99–104.
 10. Prentice FD, Wood JG. Adrenergic innervation of cat adrenal medulla. *Anat Rec*. 1975;181:689–704.
 11. Kesse WK, Parker TL, Coupland RE. The innervation of the adrenal gland. 1. The source of pre- and postganglionic nerve fibres to the rat adrenal gland. *J Anat*. 1988;157:33–41.
 12. Young JZ. The autonomic nervous system of Selachians. *Q J Microsc Sci*. 1933;75:571–624.
 13. Schober A, Wolf N, Huber K, et al. Trk B and neurotrophin-4 are important for development and maintenance of sympathetic preganglionic neurons innervating the adrenal medulla. *J Neurosci*. 1998;18(18):7272–7284.
 14. Kiss T. Experimentell- morphologische analyse der nebenniereninnervation. *Acta Anat*. 1951;XIII:81–89.
 15. Parker TL, Kesse WK, Mohamed AA, Afework M. The innervation of the mammalian adrenal gland. *J Anat*. 1993;183(2):265–276.
 16. Li C, McDonald TJ. Source of corticotropin releasing hormone-like innervation of the adrenal glands of fetal and postnatal sheep. *Brain Res*. 1997;767(1):87–91.
 17. Strack AM, Sawyer WB, Marubio LM, Loewy AD. Spinal origin of sympathetic preganglionic neurons in the rat. *Brain Res*. 1988;455:187–191.
 18. Sangari SK, Khatri K, Sengupta P. Sensory innervation of the suprarenal gland in the albino rat: a fluorescent tract tracer study. *Clin Anat*. 1998;11:1–4.
 19. Zhou XF, Oldfield BJ, Livett BG. Substance P – containing sensory neurons in the rat dorsal root ganglia innervate the adrenal medulla. *J Auton Nerv Syst*. 1991;33:247–254.
 20. Mohamed AA, Parker TL, Coupland RE. The innervation of the adrenal gland. II the source of spinal afferent nerve fibres to the guinea pig adrenal gland. *J Anat*. 1988;160:51–58.
 21. Wilson JMM, Coderre E, Renaud LP, Spanswick D. Active and passive membrane properties of rat sympathetic preganglionic neurons innervating the adrenal medulla. *J Physiol*. 2002;545(3):945–960.
 22. Kuo DC, Krauthamer GM, Yamasaki DS. The organisation of visceral sensory neurons in the thoracic dorsal root ganglia (DRG) of the cat studied by horseradish peroxidase (HRP) reaction using the cryostat. *Brain Res*. 1981;208:189–191.
 23. Neuhuber W, Niederle B. Spinal ganglion cells innervating the stomach of the rat as demonstrated by somatopetal transport of horseradish peroxidase (HRP). *Anat Embryol*. 1979;155:355–362.
 24. Cellar BG, Schramm LP. Pre- and postganglionic sympathetic activity in splanchnic nerves of rat. *Am J Physiol*. 1981;241:R55–R61.
 25. Carlsson S, Skarphedinsson JO, Jennische E, Delle M, Thoren P. Neurophysiological evidence for and characterization of the post-ganglionic innervation of the adrenal gland in the rat. *Acta Physiol Scand*. 1990;140:491–499.
 26. Hökfelt T, Lundberg JM, Schultzberg M, Fahrenkrug J. Immunohistochemical evidence for a local VIP – ergic neuron system in the adrenal gland of the rat. *Acta Physiol Scand*. 1981;113:575–576.