

radiologists should be informed of the presence of these variations and emphasize that they should report these.

92. Application of e-learning in anatomy – A knowledge, attitude and practice (KAP) study

S.K. Ghosh, S. Chakraborty, S. Biswas, S. Sharma

Department of Anatomy, ESI-PGIMS & ESIC Medical College, Joka, Kolkata, West Bengal, India

Background: This millennium has seen e-learning playing an increasingly important role in medical education as an innovative method of teaching-learning offering customized and personalized learning opportunities for students. However, perceptions of 1st year MBBS students regarding e-learning in anatomy are yet to be explored. Therefore, this KAP study was designed with both close and open ended questionnaire to identify the applications of e-learning in anatomy.

Method: A self-designed close- and open-ended questionnaire was framed to explore perceptions of 1st year MBBS students regarding applications of e-learning in studying anatomy and performance in formative assessments. Following a pilot study, the questionnaire was administered among 100 1st year MBBS students before the summative assessment. Responses were collected, analysed and recorded.

Results: All the students were found to be aware of e-learning tools available. 10% acknowledged Wikipedia as the most popular e-learning tool followed by YouTube videos (7%) and standard Google search (3%). 14% students opined that e-learning tools contributed in quick revision about anatomy, while 18% and 5% students used those for exam preparations and clarifying doubts, respectively. Overall 63% students perceived e-learning as an effective medium, while 30% students opined against it and the remaining 7% were undecided regarding its utilization while learning anatomy.

Conclusion: It can thus be concluded that majority of the first year medical students valued e-learning as an effective learning medium for anatomy. It is also suggested that e-learning in medical education curriculum would possibly be effective in improving student learning to achieve better performance related outcome.

93. Analysis of body donation in Saurashtra region: A retrospective study

V. Sharma¹, K.K. Zaveri¹, R.K. Patel¹, R.M. Patel¹, M.M. Patel¹, T.C. Singel²

¹ M. P. Shah Government Medical College, Jamnagar, Gujarat, India;

² B. J. Medical College, Ahmedabad, Gujarat, India

Introduction: Body donation is defined as the act of giving one's own body after death for medical research and education.

Aim: Aim of the present study was to analyse the pattern of body donation in Saurashtra region.

Materials and Methods: This was a retrospective study, done by collection of data through the proforma which was

obtained, at the time of body donation from the relatives and known of deceased from anatomy department of our institute, for the year 2012 and 2013.

Results: Out of total 40 cases taken, 95% were above 50 years, 75% were male, 67% were literate and 63% were non-working.

Conclusion: Hence concluded that the most of the body donated to our institute were aged >50 years, most of them were literate and were non-working at the time of death. Males were found to outnumber females in body-donation.

94. Olanzapine induced placental changes in mice

Anand Mishra, Soumya Khanna

Department of Anatomy, IMS, BHU, India

Introduction: Olanzapine, a typical antipsychotic is mainly used in treatment of schizophrenia and bipolar disease.

Materials & Methods: Olanzapine was given to pregnant mice in doses of 0.2 mg/kg and 2 mg/kg on day 6–12 of gestation, while control mice were given distilled water on same days of gestation. The mice were sacrificed on day 19th of pregnancy by deep ether anaesthesia, and placentae were collected after performing uterotomy. The placenta were weighed, observed for overt anomaly, fixed 10% formalin, and processed for histological study by staining with H&E.

Results: The treated placenta shows a dose dependent hyalinization and thickening of trichorial membrane and disruption of maternal venous sinusoid.

Conclusion: Olanzapine causes toxicity in placenta and thus can harm the fetus so it should be used with caution in pregnancy.

95. Effect of consanguinity on congenital defects

Charmode Sundip Hemant

Esic Medical College Parippally, Kollam, Kerela, India

Objective: To determine the Influence of Consanguinity in occurrence of congenital anomalies and the occurrence of more common types of congenital anomalies.

Methods: The study was done retrospectively in Government medical college and its attached hospitals during the period of 1st October 2010 till 31st May 2012. A total of 182 congenitally anomalous live births and still births admitted in above hospitals during the same period were studied against total deliveries taken place (10,114). Their parents were enquired using Questionnaires and information regarding consanguinity, degree of consanguinity, type and subtype of anomaly etc. was obtained. Observations were statistically analysed and compared with previous studies.

Results: Occurrence of congenital anomalies was 1.16%. Cardiovascular anomalies (39 cases) were the most common type of malformation and Atrial septal defect (18 cases) the most common subtype. Most common form of Degree of consanguinity found among consanguineously married parents with anomalous births was third degree/first cousins, half uncles

and half niece, and half aunt and half nephew. Most of the births were male and first order. Male:Female ratio was 1:2.3. Frequency of consanguineous marriages was found more in Muslims than Hindus (statistically significant).

Conclusion: Congenital malformations were found to occur more common in consanguineous unions than non-consanguineous unions (Statistically insignificant). Cardiovascular anomalies were the most common type of malformation and atrial septal defect, the most common subtype. Consanguinity was found more common to occur in muslim population than hindus.

96. Development of parafollicular cells and its relationship with developing thyroid follicles in human fetuses

S.S. Das, J.M. Kaul, S. Mishra

Maulana Azad Medical College, New Delhi, India

Objective: The parafollicular cells or C (clear) cells in man is a part of neuroendocrine system, classified under "amine precursor uptake and decarboxylation" (APUD) cells. Their role in adults has been *reputable* but in fetus is still unclear. The present study was a baseline study endeavoring to describe the chronological development of the parafollicular cells with particular focus on its correlation with developing human thyroid follicles.

Methods: The study was conducted on 10 aborted fetuses (14-28 weeks), procured from the Department of Obstetrics and Gynaecology, Lok Nayak Hospital, New Delhi. Serial sections of fetal thyroid gland were generated, stained with haematoxylin and eosin, Sevier Munger stain, and immunohistochemistry using the anti- calcitonin antibody and examined qualitatively.

Results: In our study, the parafollicular cells were seen as earlier as by 14th week. They became morphologically and functionally mature by 16th week of gestation. The parafollicular cells were getting organized from scattering to parafollicular location then to a more localized area, i.e., intrafollicular along with the follicular development. As the follicles were enlarging, the intrafollicularly located parafollicular cells which were initially present in groups were getting displaced singly between the follicular cells in the same follicle.

Conclusion: The sequential development pattern of the parafollicular cells in relation to developing thyroid follicles was established. This immunohistochemical study also concluded that the parafollicular cells might have higher character to play in the early gestational age such as regulation of ossification in the human fetus.

97. A morphological and histological study of placentae in respect of hypertensive gravid mother

Sutandro Choudhury, Sitansu Panda,
Prafulla Kumar Chinara

I.M.S. & SUM Hospital, Bhubaneswar, Odisha, India

Objective: A comparative study of placentae of the normotensive and hypertensive mothers was undertaken as perinatal death is a serious public health problem in our country. Many of such deaths are due to birth of preterm babies and hypertension of the gravid mother is one of the high risk factor for the same.

Methods: Altogether 60 placentae of which 30 placentae were from normotensive mother and 30 placentae from hypertensive (both pre-existing and pregnancy induced) were collected from the Department of Obstetrics & Gynaecology of I.M.S. & SUM Hospital, Bhubaneswar, Odisha. Study was carried out in the Department of Anatomy of the same Institute. The morphological and morphometrical parameters of both the groups were compared. Care has been taken to note the areas of infarction and calcification. Histological changes were also noted.

Results: Placentae from hypertensive gravid mothers were found to be lesser in weight, surface area, thickness and volume. Greater number of areas of calcification and areas of infarction were noted in the study group. Histological study revealed an increase in syncytial knot count and decrease in the percentage of villi.

Conclusion: Hypertension of the gravid mothers is responsible for the changes as specified and corroborate with the findings of the previous studies.

98. The histomorphometry of islets in the mammalian pancreas

S.G. Deka¹, K.L. Talukdar², J. Sarma², T. Sarma¹

¹Fakhruddin Ali Ahmed Medical College, Barpeta, India; ²Gauhati Medical College, Guwahati, India

A study on the histology of the pancreas is undertaken to observe and compare the islets of five different mammals. The main objective of this study is to ascertain possible histomorphometrical closeness of the animal (mammal) pancreatic islet with the human pancreatic islet, so as to explore potential xenogenic sources of donors for organ transplantation. The study was conducted on five mammals – the rat, the rabbit, the goat, the pig and the human being. The pancreas was sectioned and stained for light microscopy by Haematoxylin and Eosin. Well formed islets, acini and ducts were present in all the mammalian pancreases. The islets appeared as light stained encapsulated areas. The islets were observed for their shape, staining character, location, whereas its dimensions were measured with the help of a micrometer slide used with the light microscope. The distribution of islets within the pancreas was another aspect of the study. The biometrical values of different groups were statistically analyzed according to Croxton. The study showed that in the five mammalian groups, variations in length, breadth, diameter, volume of the islets had no significant statistical difference, despite the fact that the mammals had wide differences in average body weights and pancreatic weights.