

14

Morphological variations of musculocutaneous and median nerves in arm



Stuti Tandon^{1,*}, P. Singh¹, R.S. Rajpoot², Rani Archana¹, A.K. Srivastava¹

¹ King George's Medical University, Lucknow, India

² RIMSR, Saifai, Etawah, Uttar Pradesh, India

Aims and objectives: The study was aimed to observe variations in the branching pattern of musculocutaneous and median nerves.

Material and methods: The present study was performed by routine dissections on 38 cadavers at King George's Medical University UP, Lucknow.

Results: Out of 38 cases studied, in 6 cases were observed in which variation of musculocutaneous and median nerves were seen. In one case, musculocutaneous nerve on left side arose as a slender branch from the lateral cord, supplied and pierced coracobrachialis muscle and ended. In the same case just after its formation, median nerve gave a branch which descended between biceps and brachialis, sending branches to both and emerged from beneath the lateral border of tendon of biceps as the lateral cutaneous nerve of forearm. In three limbs there were longer roots of the median nerve which united at the level of mid arm and were intercommunicating. In another left upper limb, apart from long roots, after formation of median nerve a branch was given off which descended downwards and laterally and communicated with musculocutaneous nerve at the level of elbow joint. In another limb there was a communication between the median nerve and the musculocutaneous nerve at the level of mid arm.

Conclusion: Such variations are important clinically for surgeons performing surgeries on upper limb.

Conflicts of interest

The authors have none to declare.

<http://dx.doi.org/10.1016/j.jasi.2016.08.022>

15

Fetal foot and hand lengths: Their relationship with crown-rump (CR) length and gestational age



Garima Sharma*, N.S. Thounaojam

Department of Anatomy, Regional Institute of Medical Sciences, Imphal, Manipur, India

Aims and objectives: To study the relationship between fetal crown rump length and fetal hand and foot length, thereby determining the accuracy in estimating gestational age by a cross sectional study.

Material and methods: A total of 100 formalin fixed fetuses of Manipuri origin, obtained from the Department of Obstetrics and Gynaecology, Regional Institute of Medical Sciences, Imphal were studied in the Department of Anatomy from February to July 2015. The parameters measured in fetuses were: (i) CR length, (ii) foot length, and (iii) hand length. The data was analysed using SPSS software by regression analysis. Graphs were also plotted to determine pattern of growth and their correlation with crown rump length, if any.

Results: Out of 100 fetuses (43 female and 57 male) the mean foot length and hand length progressively increased with increase in crown rump length. Measurements were not significantly different on right or left side or among male and female fetuses.

A statistically significant linear relationship was found between foot length and crown rump length ($r = 0.980$, $p < 0.0001$) and hand length and crown rump length of the fetus ($r = 0.986$, $p < 0.0001$).

Conclusion: In the present study fetal hand and foot lengths have been found to highly correlate with crown rump length and thereby gestational age. Therefore these parameters could be utilized to estimate gestational age.

Conflicts of interest

The authors have none to declare.

<http://dx.doi.org/10.1016/j.jasi.2016.08.023>

16

A study of highest nuchal line in north Indian crania



S.H.H. Zaidi

Rohilkhand Medical College, 243006 Uttar Pradesh, India

Aims and objectives: Studies of non-metric cranial variants have been a field of considerable interest to research workers especially because of their racial and regional importance.

Material and methods: 18 north Indian skulls of UP were studied for the highest nuchal line a cranial variant in the present study.

Results: Highest nuchal line variant was seen in 50% cases.

Conclusion: Findings are discussed and compared with other global studies and are found to be of considerable regional and racial significance.

Conflicts of interest

The author has none to declare.

<http://dx.doi.org/10.1016/j.jasi.2016.08.024>

17

Morphometrical study of thyroid cartilage



Patel Shital, Bhardwaj Rashmi*, V.H. Vaniya

Medical College, Baroda, Gujarat, India

Aims and objectives: To find out dimension of various parameters of thyroid cartilage, variation, differences among all the data and to compare data between males and females.

Material and methods: Thirty samples of adult thyroid cartilage (16 males and 14 females) of known sex were taken from embalmed cadavers. Morphometric measurements were taken using a digital vernier caliper and optical goniometer to see normal variation, sex differences and bilateral asymmetry. Observations were analysed statistically.

Results: Parameters measured in thyroid cartilage were height (average in males 31.19 mm and in females 24.17 mm), width (average in males 39.73 mm and in females 30.00 mm), posterior border length (average in males 45.31 mm and in females 35.40 mm) of thyroid lamina and thyroid angle (average in males 85.75 and in females 98.86), show significant differences in males and females. All the parameters of thyroid cartilage except thyroid angle were higher in males than females. Bilateral asymmetry in the size of thyroid lamina was seen in majority of specimens. However, in 3 cases (10%) thyroid prominence was not in median plane, deviated towards left side. In one case asymmetry was also associated with asymmetry in direction of superior cornu.