

Conflicts of interest

The authors have none to declare.

<https://doi.org/10.1016/j.jasi.2018.06.191>

38

Accessory sulci and segmentation on the surface of the liver: a clinical pitfall



Chetan Sahni*, Vishwajit Ravindra Deshmukh, Suryakanta Seth

All India Institute of Medical Sciences, New Delhi, India

Introduction: Diaphragmatic surface of liver is usually smooth but sometimes it is well marked by the sulci or indentations. Sometimes these sulci were so deep to be termed as fissures as they divide the lobe of liver into different segments. We report the case of accessory sulci along the anterosuperior surface of the right lobe associated with Reidel's lobe along the inferior border of the left lobe of same specimen. In another specimen, the sulci were so deep to be termed as fissures, which divides the caudate and quadrate lobe. Knowledge about the accessory sulci and lobes were necessary as they may appear as incidental finding during the laparoscopic examinations. Hence, it is very much promising to know about these sulci and lobes for hepatobiliary surgeons, anatomists and radiologists.

Observation/Result: During routine dissection classes conducted for first-year MBBS students in the Department of Anatomy, All India Institute of Medical Sciences, New Delhi, India, we encountered the presence of indentation or furrows on the anterosuperior surface of the right lobe of the liver in male cadaver aged 60 years and the division of caudate and quadrate lobe in female cadaver aged 55 years.

Discussion: Accessory lobe in present case report is Riedel's lobe in one of the cadaver, is clinically very important lobe resulting in misdiagnosis. Hence, it should always be considered in patients undergoing the cross-sectional studies. It may be a port of disease that might not be demonstrated until and unless the inferior border of the liver was examined.

Conflicts of interest

The authors have none to declare.

<https://doi.org/10.1016/j.jasi.2018.06.192>

39

Insular cortex: variation in pattern of gyri with interhemispheric difference



JayaKadam Deshmukh*, Vandana Sharma, Sonia Singh Baweja

Department of Anatomy, Gandhi Medical College, Bhopal, India

Background and aims: Variations in the morphological pattern of gyri of insular cortex is very common. Usually insular cortex represents 5 to 7 gyri, sometimes with bifid/trifid or hypoplastic gyrus. During the routine thesis related dissection we got a rare variant having hypoplastic gyrus on right side and trifid gyrus on left side.

Materials and methods: The study includes 20 formalin fixed human cadaveric brains (40 cerebral hemispheres) in the Anatomy Department, Gandhi Medical College, Bhopal. Measurements were

taken with the help of digital vernier caliper (in mm with 2 decimal), thread, artery forceps. We measured the insular cortex peripheral sulci (anterior, posterior, superior, inferior) and central insular sulci. Then all 7 gyri—transverse gyrus (TG), accessory (ACG), anterior short gyrus (ASG), middle short gyrus (MSG), posterior short gyrus (PSG), anterior long gyrus (ALG), posterior long gyrus (PLG) lengths and pattern (absent gyrus, bifid, trifid, hypoplastic) were studied and recorded.

Results: common variant with all 7 gyri on both sides. The rare variant represents with hypoplastic MSG on right side and bifid ASG on both side and trifid PSG in left side cerebral hemisphere.

Conclusion: The insula shows unique pattern in every individual, having significant interhemispheric variation and it is also declared morphologically most variable structure of brain with great clinical and neurosurgical significance. Trans insular and trans sylvian approach is very common in neurosurgical assess and it is essential to highlight every variable and interhemispheric difference as we studied through our neuroanatomical parameters observational records.

Conflicts of interest

The authors have none to declare.

<https://doi.org/10.1016/j.jasi.2018.06.193>

40

Anomalous branching pattern from aortic arch: a case report



Tanveer Ahmad

Department of Anatomy, Faculty of Dentistry, Jamia Millia Islamia, New Delhi, India

Increasing in the fields of cardiac and vascular surgery has served to revive interest in the developmental and adult anatomy of the aortic arches and the great vessels derived from the area. Under normal circumstances, three branches originate from the aortic arch: the brachiocephalic trunk, left common carotid artery, and left subclavian artery, in about 80% of individuals. Branches originating from the aortic arch may show different configurations. As far as the branches of the aortic arch are concerned, there are numerous possible variations in the origins of these branches. The anatomic and morphologic variations of the aortic arch branches are important for diagnostic. The knowledge of the branching pattern of aortic arch is important during supra-aortic angiography, aortic instrumentation, thoracic and neck surgery. The purpose of this study is to describe different branching patterns of arch of aorta in order to offer useful data to anatomists, radiologists, vascular surgeons doing surgical procedures in the head and neck region while relating it to the embryological basis.

In the present study, left internal carotid artery, left external carotid artery and left subclavian artery had an aberrant origin, however, the anatomical courses of these arteries were found normal. Details will be discussed in full length paper.

An abnormal origin of the left common carotid artery from the initial portion of the brachiocephalic trunk was found in the superior mediastinum in a 81-year-old Caucasian male cadaver during dissection practice. We report on the exact morphology of that variant that is appeared in an incidence of 0.2% in the literature. We discuss the relative literature and pay attention on the significance of such a variation for clinicians in its recognition and protection.

Keywords: developmental anatomy, variations, vascular surgery.